

# Bitter Gourd Cultivation

## Climate and Soil Requirements

Bitter gourd thrives in warm and humid climates. The optimal temperature range for cultivation lies between 25°C and 35°C. Adequate exposure to full sunlight is essential for its growth and fruit development. While bitter gourd can flourish in a variety of soils, well-drained loamy soil enriched with organic matter is preferable. The pH level of the soil should range between 6.0 and 7.0 to promote optimal growth. Bitter gourd is propagated through seeds, which can be directly sown in the field or nurtured in nurseries for later transplantation. Soak the seeds in water for 24 hours before planting to enhance germination rates. Space the seeds at intervals of 1.5 to 2 meters between rows and 1 meter between individual plants.

## Land Preparation

Begin by plowing and harrowing the land to create a fine tilth. Incorporate well-decomposed organic matter or farmyard manure into the soil prior to planting to enhance soil fertility and structure. To ensure effective drainage, establish raised beds or ridges.

## Planting

In tropical climates, bitter gourd can be cultivated year-round. For temperate regions, it is advisable to plant after the last frost has passed and the soil has warmed. Transplant seedlings that are three to four weeks old into prepared beds or ridges, being cautious not to damage the roots during transplantation.

## Irrigation and Support

Regular watering is crucial, particularly during dry spells.

Maintain consistent intervals of irrigation to keep the soil adequately moist without causing waterlogging. Drip irrigation is recommended for efficient water utilization and to prevent fungal diseases. Bitter melon is a vine crop that benefits from trellising or staking. Install sturdy trellises or stakes during planting to support the growing vines. Guide the vines to climb the trellises or stakes, making fruit harvesting and disease management more convenient.

## **Fertilization**

Apply a balanced fertilizer during land preparation or transplanting. After the first month of growth, side-dress the plants with nitrogen-rich fertilizer. Monitor the crop's nutrient requirements and administer additional fertilizer as needed.

## **Weed and Pest Management**

Maintain a weed-free environment through regular weeding or mulching. Regularly inspect the crop for pest infestations, including aphids, fruit flies, and spider mites. Employ organic or chemical pesticides as necessary, adhering to recommended dosages and safety precautions.

**Harvesting:** Bitter melon fruits are typically ready for harvest around 60 to 70 days after transplantation. Harvest the fruits when they are green, firm, and not overripe. Utilize a sharp knife or pruning shears to gently cut the fruits from the vine, taking care to avoid bruising or damage.

## **Post-Harvesting and Storage**

After harvesting, sort and grade the fruits based on size and quality, discarding any damaged or diseased ones. Bitter melon can be stored for a short duration in a cool, dry place or refrigerated to extend its shelf life. Successful bitter melon farming entails consistent monitoring, timely pest and disease management, proper irrigation, and appropriate fertilization

practices to ensure a productive and healthy crop.